



Generalized connected domination in graphs

Mekkia Kouider, Preben Dahl Vestergaard

► To cite this version:

Mekkia Kouider, Preben Dahl Vestergaard. Generalized connected domination in graphs. Discrete Mathematics and Theoretical Computer Science, 2006, Vol. 8, pp.57-64. 10.46298/dmtcs.373 . hal-00961101

HAL Id: hal-00961101

<https://inria.hal.science/hal-00961101>

Submitted on 19 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Generalized connected domination in graphs

Mekkia Kouider¹ and Preben Dahl Vestergaard^{2†}

¹*Laboratoire de Recherche en Informatique, UMR 8623, Bât. 490, Université Paris Sud, 91405 Orsay, France.*

E-mail: km@lri.fr

²*Dept. of Mathematics, Aalborg University, Fredrik Bajers Vej 7G, DK-9220 Aalborg Ø, Denmark.*

E-mail: pdv@math.aau.dk

received Nov 10, 2003, revised Oct 19, 2005, accepted Jan, 2006.

As a generalization of connected domination in a graph G we consider domination by sets having at most k components. The order $\gamma_c^k(G)$ of such a smallest set we relate to $\gamma_c(G)$, the order of a smallest connected dominating set. For a tree T we give bounds on $\gamma_c^k(T)$ in terms of minimum valency and diameter. For trees the inequality $\gamma_c^k(T) \leq n - k - 1$ is known to hold, we determine the class of trees, for which equality holds.

Keywords: connected domination, domination, tree

Mathematics Subject Classification: 05C69

1 Introduction

We consider simple non-oriented graphs. The largest valency in G is denoted by $\Delta(G) = \Delta$, the smallest by $\delta(G) = \delta$. By P_n we denote a path on n vertices and C_n denotes a circuit on n vertices. In a graph a **leaf** or **pendant vertex** is a vertex of valency one and a **stem** is a vertex adjacent to at least one leaf. In K_2 each vertex is both a leaf and a stem. The set of leaves in a graph G is denoted by $\Omega(G)$. The set of neighbours to a vertex x is denoted $N(x)$. By $K_{1,k}$ we denote a star with one central vertex joined to k other vertices. A **subdivided star** is a star with a subdivision vertex on each edge. By the **corona graph** on H we understand the graph $G = H \circ K_1$ obtained from the graph H by adding for each vertex x in H one new vertex x' and one new edge xx' . In a corona graph each vertex is either a leaf or a stem adjacent to exactly one leaf. In particular, if H is a tree, we obtain a **corona tree** $T = H \circ K_1$.

The **eccentricity** $e(x)$ of a vertex x is defined by $e(x) = \max\{d(x, y) | y \in V(G)\}$. The **diameter** of G is $\text{diam}(G) = \max\{e(x) | x \in V(G)\}$. Let $D \subseteq V(G)$, then $N(D)$ is the set of vertices which have a neighbour in D and $N[D]$ is the set of vertices which are in D or have a neighbour in D , $N[D] = D \cup N(D)$. A set $D \subseteq V(G)$ **dominates** G if $V(G) \subseteq N[D]$, i.e. each vertex not in D is adjacent to a vertex in D . The **domination number** $\gamma(G)$ is the cardinality of a smallest dominating set in G .

For a given graph G it is NP-hard to determine its domination number $\gamma(G)$, but we can search for for upper bounds as O. Ore started doing about fifty years ago. Also it may be more tractable to restrict the

[†]Supported by the Danish Natural Science Research Council

minimum dominating set problem to consider only such dominating sets which induce a connected subset of G , this problem is called **the minimum connected dominating problem** and it is still NP-complete; In network design theory it is called the **maximum leaf spanning tree problem** [4], the name will be clear from Section 2 below. We shall study a concept intermediate to the classical and the connected domination, namely by demanding the dominating set to induce at most a given number k of components, we aim at presenting upper bounds for its order γ_c^k . Quite likely there is a corresponding problem in network design theory, although we are aware of no reference.

A comprehensive introduction to domination theory is given in [7, 14] and variations are discussed in [5, 13, 15].

Ore [10] proved the inequality below while C. Payan and N. H. Xuong [11], Fink, Jacobsen, Kinch and Roberts [3] determined its extremal graphs.

Proposition 1 *Let G be a connected graph with n vertices, $n \geq 2$. Then $\gamma(G) \leq \frac{n}{2}$ and equality holds if and only if G is either a corona graph or a 4-circuit.*

If a tree T has $\gamma(T) = \frac{n}{2}$, then n is even and Proposition 1 implies that T is a corona tree.

Definition For a positive integer k and a graph G with at most k components we define

$$\gamma_c^k(G) = \min \{|D| \mid D \subseteq V(G), D \text{ has at most } k \text{ components and } D \text{ dominates } G\}.$$

A set D attaining the minimum above is called a γ_c^k -set for G .

Example

$$\gamma_c^k(P_n) = \gamma_c^k(C_n) = \begin{cases} n - 2k & \text{for } n \geq 3k \\ \lceil \frac{n}{3} \rceil & \text{for } 1 \leq n \leq 3k \end{cases}$$

For $k = 1$ we have that γ_c^1 is the usual connected domination number, $\gamma_c^1(G) = \gamma_c(G)$.

There exists for every graph G a k such that $\gamma_c^k(G) = \gamma(G)$, e.g. $k = |G|$.

For G connected and $k \geq 1$, obviously, $\gamma(G) \leq \gamma_c^k(G) \leq \gamma_c(G)$.

2 General graphs

Let G be a connected graph with n vertices and k a positive integer. Let $\epsilon_F(G)$ be the maximum number of leaves among all spanning forests of G , and $\epsilon_T(G)$ be the maximum number of leaves among all spanning trees of G . With this notation Niemen [9] proved statement (i) below about γ and Hedetniemi and Laskar [8] generalized it to statement (ii) about γ_c .

$$(i) \quad \gamma(G) = n - \epsilon_F(G),$$

$$(ii) \quad \gamma_c(G) = n - \epsilon_T(G).$$

In the next two theorems we extend these results to γ_c^k .

Theorem 1 *Let G be a connected graph with n vertices and k a positive integer. Let $\epsilon_{F_k}(G)$ be the maximum number of leaves among all spanning forests of G with at most k trees. Then*

$$\gamma_c^k(G) = n - \epsilon_{F_k}(G).$$

Proof: In any spanning forest F with at most k trees the leaves will be dominated by their stems, so $\gamma_c^k(G) \leq n - |\Omega(F)|$ and hence $\gamma_c^k(G) \leq n - \epsilon_{F_k}(G)$.

Conversely, let $D = D_1 \cup D_2 \cup \dots \cup D_t$, $1 \leq t \leq k$, be a γ_c^k -set for G . Choose for each D_i a spanning tree T_i , $1 \leq i \leq t$. For each vertex in $V(G) \setminus D$ choose one edge which is incident with a vertex in D . We have constructed a spanning forest F with t components and at least $n - |D| = n - \gamma_c^k(G)$ leaves. Therefore $\epsilon_{F_k}(G) \geq n - \gamma_c^k(G)$ and Theorem 1 is proved. \square

Theorem 2 Let k be a positive integer and G a connected graph. Then

$$\begin{aligned} \gamma_c^k(G) &= \min \{ \gamma_c^k(F_k) \mid F_k \text{ is a spanning forest of } G \text{ with at most } k \text{ trees} \} \\ &= \min \{ \gamma_c^k(T) \mid T \text{ is a spanning tree of } G \} \end{aligned}$$

Proof: Let F_k be a spanning forest of G with at most k trees. Certainly $\gamma_c^k(G) \leq \gamma_c^k(F_k)$ since a set which dominates F_k also dominates G . Conversely, we can in G find a spanning forest F_k with at most k components such that $\gamma_c^k(G) = \gamma_c^k(F_k)$: As was originally also done in the proofs for (i) and (ii) above we construct F_k from a γ_c^k -set $D = D_1 \cup D_2 \cup \dots \cup D_t$, $1 \leq t \leq k$, by choosing a spanning tree T_i in each connected subgraph D_i and joining each vertex in $V(G) \setminus D$ to precisely one vertex in D . Obviously, $\gamma_c^k(F_k) \leq |D| = \gamma_c^k(G)$. This proves the first equality. For the second equality we observe that the first minimum is chosen among a larger set, so that $\min \gamma_c^k(F_k) \leq \min \gamma_c^k(T)$, and also that any F_k by addition of edges can produce a tree T with $\gamma_c^k(T) \leq \gamma_c^k(F_k)$. \square

Hartnell and Vestergaard [6] proved the following result.

Proposition 2 For $k \geq 1$ and G connected

$$\gamma_c(G) - 2(k - 1) \leq \gamma_c^k(G) \leq \gamma_c(G).$$

From Proposition 2 we can easily derive the following corollary which is a classical result proven by Duchet and Meyniel. [2]

Corollary 3 For any connected graph G , $\gamma_c(G) \leq 3\gamma(G) - 2$.

Proof: Let G be a connected graph with domination number $\gamma(G)$. Choose $k = \gamma(G)$, then $\gamma_c^k(G) = \gamma(G)$. Substituting into Proposition 2 above we obtain $\gamma_c(G) - 2(k - 1) \leq \gamma(G)$ and that proves the corollary. \square

2.1 Other bounds on γ_c^k

Theorem 4 For a positive integer k and a connected graph G with maximum valency Δ we have

(A) $\gamma_c(G) \leq n - \Delta$ and for trees T equality holds if and only if T has at most one vertex of valency ≥ 3 .

(B) $\gamma_c^k(G) \leq n - \frac{(r-1)(\delta-2)}{3} - 2k$ if G has diameter $r \geq 3k-1$ and the minimum valency $\delta = \delta(G)$ is at least 3.

(C) If G is a connected graph with two vertices of valency Δ at distance d apart, $d \geq 3$, then

$$\gamma_c^k(G) \leq n - 2(\Delta - 1) - 2 \min\{k - 1, \frac{d - 2}{3}\}. \quad (1)$$

(D) Let $x \in V(G)$ have valency $d(x)$ and eccentricity $e(x)$. Then

$$\gamma_c^k(G) \leq n - d(x) - 2 \min\{k - 1, \frac{e(x) - 2}{3}\}. \quad (2)$$

Proof:

(A) Let T be a spanning tree of G with $\Delta(T) = \Delta(G) = \Delta$, then T has at least Δ leaves, and hence $\gamma_c(G) \leq \gamma_c(T) \leq n - \Delta$.

If T has two vertices of valency ≥ 3 , the number of leaves in T will be larger than Δ , and we get strict inequality in (A). Clearly, a tree T with exactly one vertex of valency $\Delta \geq 3$ has equality in (A) and for $\Delta = 2$, we obtain a path P_n with $\gamma_c(P_n) = n - 2$.

(B) Let $P = v_1 v_2 v_3 \dots v_{3t+u}$, $k \leq t, 0 \leq u \leq 2$, be a diametrical path in G . The diameter of T equals the length of P , which is $r = 3t + u - 1$. For $i = 1, \dots, t$ let v_{3i-1} have neighbours v_{3i-2}, v_{3i} on P and a_{ij} off P , $j = 1, \dots, s_i$, $s_i \geq \delta - 2 \geq 1$. In $G - \{v_{3i} v_{3i+1} | 1 \leq i \leq k - 1\}$ consider the $k - 1$ disjoint stars with center v_{3i-1} and neighbours $N(v_{3i-1})$, $1 \leq i \leq k - 1$, and the remaining tree to the right consisting of the path $v_{3k-2} v_{3k-1} v_{3k} \dots v_{3t+u}$ and leaves $v_{3i-1} a_{3i-1}$, $j = 1, \dots, s_i$, $s_i \geq \delta - 2 \geq 1$ adjacent to vertices v_{3i-1} , $k \leq i \leq t$.

Extend this forest of k trees to a spanning forest F with k trees in $G - \{v_{3i} v_{3i+1} | 1 \leq i \leq k - 1\}$. The number of leaves in F is at least $t(\delta - 2) + 2k$ and hence $\gamma_c^k(G) \leq n - t(\delta - 2) - 2k$. From

$$t = \frac{r + 1 - u}{3} \geq \frac{r - 1}{3} \text{ we obtain the desired result } \gamma_c^k(G) \leq n - \frac{(r - 1)(\delta - 2)}{3} - 2k.$$

C Let v_1, v_s be two vertices in G with maximum valency, $d(v_1) = d(v_s) = \Delta$, and let $P = v_1 v_2 \dots v_s$ be a shortest $v_1 v_s$ -path, $s = 3t + 1 + u$, $t \geq 1, 0 \leq u \leq 2$.

Case 1, $t \geq k - 1$: In $G - \{v_{3i-1} v_{3i} | 1 \leq i \leq k - 2\}$ we extend the k trees listed below to a spanning forest F of G ,

1. The star consisting of v_1 joined to all its neighbours,
2. the $k - 2$ paths of length two $v_{3i} v_{3i+1} v_{3i+2}$, $1 \leq i \leq k - 2$,
3. the path $v_{3k-3} v_{3k-2} \dots v_s$ together with all $\Delta - 1$ neighbours of v_s outside of P .

F will have at least $2(\Delta - 1) + 2(k - 1)$ leaves.

Case 2, $t \leq k - 2$: $s = 3t + 1 + u$, $d = d(v_1, v_s) = s - 1 = 3t + u$, $t - 1 = \frac{d - u}{3} - 1 \geq$

$\frac{d - 2}{3} - 1$. As before, we can find a spanning forest F of G whose number of leaves is at least

$$2\Delta + 2(t - 1) \geq 2(\Delta - 1) + 2\frac{d - 2}{3} \text{ and consequently } \gamma_c^k(G) \leq n - 2(\Delta - 1) - 2\frac{d - 2}{3}.$$

The proof of D is similar.

□

3 Trees

For trees Hartnell and Vestergaard [6] found

Proposition 3 *Let k be a positive integer and T a tree with $|V(T)| = n, n \geq 2k + 1$. Then $\gamma_c^k(T) \leq n - k - 1$.*

This inequality is best possible. For $k = 1$ the extremal trees are paths P_n and for $k \geq 2$ extremal trees will be described in the following Theorem 5.

A tree T is of type A if it contains a vertex x_0 such that $T - x_0$ is a forest of trees $T_1, T_2, \dots, T_\alpha, \alpha \geq 1$, such that each tree T_i is a corona tree and x_0 is joined to a stem in each of the trees $T_i, 1 \leq i \leq \alpha$. We note that a subdivision of a star is a tree of type A.

A tree T is of type B if it contains a path uvw such that $T - \{u, v, w\}$ is a forest of corona trees $T_1, T_2, \dots, T_s, T_{s+1}, \dots, T_\alpha, \alpha \geq 2, 1 \leq s < \alpha$ and u is joined to a stem in each of the trees T_1, T_2, \dots, T_s , while w is joined to a stem in each of the trees T_{s+1}, \dots, T_α .

Proposition 4 below was proven by Randerath and Volkmann [12], Baogen, Cockayne, Haynes, Hedetniemi and Shangchao [1].

Proposition 4 *If T is a tree with n vertices, n odd, and $\gamma(T) = \lfloor \frac{n}{2} \rfloor$ then T is a tree of type A or B.*

We shall now determine the trees extremal for Proposition 3.

Theorem 5 *Let $k \geq 2$ be a positive integer and T a tree with n vertices, $n \geq 2k + 1$. Then $\gamma_c^k(T) = n - k - 1$ if and only if one of cases (i)-(iii) below occur:*

- (i) $k = \frac{n-1}{2}, \gamma_c^k(T) = \gamma(T) = \frac{n-1}{2}$ and T is of type A or B.
- (ii) $k = \frac{n-2}{2}, \gamma_c^k(T) = \gamma(T) = \frac{n}{2}$ and T is a corona tree.
- (iii) $k = \frac{n-3}{2}, \gamma_c^k(T) = \frac{n+1}{2}, \gamma(T) = \frac{n-1}{2}$ and T is a star $K_{1,k+1}$ with a subdivision vertex on each edge.

Proof: First, let $k \geq 2$ and a tree T of order n be given such that $n \geq 2k + 1$ and $\gamma_c^k(T) = n - k - 1$. We shall prove that T is as described in one of the three cases (i)-(iii).

We note in passing that

Remark 1 $\gamma(T) \leq k$ implies $\gamma_c^k(T) = \gamma(T)$, and that likewise $\gamma_c^k(T) \leq k$ implies $\gamma_c^k(T) = \gamma(T)$.

If $n = 2k + 1$, or equivalently $k = \frac{n-1}{2}$, we have by assumption $\gamma_c^k(T) = n - k - 1 = k$ and, as just observed above, that implies that also $\gamma(T) = k$. Since $k = \lfloor \frac{n}{2} \rfloor$ we obtain from Proposition 4 that T is a tree of type A or B, so Case (i) occurs.

If $n = 2k + 2$, or equivalently $k = \frac{n-2}{2}$ we have by assumption $\gamma_c^k(T) = n - k - 1 = k + 1$. Certainly $\gamma(T) \leq \gamma_c^k(T)$, but if $\gamma(T) \leq k$ then we should have that $\gamma_c^k(T) = \gamma(T) \leq k$ in contradiction

to $\gamma_c^k(T) = k + 1$, therefore $\gamma(T) = k + 1 = \frac{n}{2}$. From Proposition 1 we obtain that T is a corona tree, i.e. Case (ii) occurs.

We may now assume $n \geq 2k + 3$, and we shall prove that, in fact, n equals $2k + 3$ and that Case (iii) occurs.

Let $v_1 v_2 \dots v_\alpha$ be a longest path in T . Since $\gamma_c^k(T) = n - k - 1 \geq k + 2 \geq 4$, T is neither a star nor a bistar and therefore $\alpha \geq 5$. We must have $d_T(v_2) = 2$, because otherwise $d_T(v_2) \geq 3$ and we could from T delete three leaves adjacent to v_2 , if $d_T(v_2) \geq 4$, and in case $d_T(v_2) = 3$ we could delete v_2 and its two adjacent leaves. In both cases we would obtain a tree T' of order $n - 3 \geq 2(k - 1) + 1$ which by Proposition 3 has $\gamma_c^{k-1}(T') \leq (n - 3) - (k - 1) - 1 \leq n - k - 3$. Adding v_2 to a $\gamma_c^{k-1}(T')$ -set we would obtain $\gamma_c^k(T) \leq n - k - 2$, a contradiction. Therefore $d_T(v_2) = 2$.

The vertex v_3 cannot be adjacent to two leaves c and d , say, because, then the tree $T' = T - \{v_1, v_2, c, d\}$ would have order $n - 4 \geq 2(k - 1) + 1$. Thus Proposition 3 gives that $\gamma_c^{k-1}(T') \leq (n - 4) - (k - 1) - 1 \leq n - k - 4$ and adding v_2, v_3 to a $\gamma_c^{k-1}(T')$ -set we would obtain $\gamma_c^k(T) \leq n - k - 2$, a contradiction. So v_3 can be adjacent to at most one leaf. The case $d_T(v_3) = 3$ and v_3 adjacent to one leaf c can similarly be seen to be impossible by considering $T' = T \setminus \{v_1, v_2, v_3, c\}$.

On the other hand $d_T(v_3) \geq 3$, for assume $d_T(v_3) = 2$, then $T' = T \setminus \{v_1, v_2, v_3\}$ has $\gamma_c^{k-1}(T') \leq n - k - 3$ and addition of v_2 to a $\gamma_c^{k-1}(T')$ -set would give $\gamma_c^k(T) \leq n - k - 2$, a contradiction.

Assume therefore that v_3 besides v_2 and v_4 is adjacent to precisely one leaf c and to at least one further vertex a , where a has valency two and is adjacent to the leaf b . Then $T' = T \setminus \{v_1, v_2, a, b\}$ has order $n - 4 \geq 2(k - 1) + 1$ and Proposition 3 gives that (3) $\gamma_c^{k-1}(T') \leq (n - 4) - (k - 1) - 1 \leq n - k - 4$. In T' the vertex c is a leaf and as any γ_c^{k-1} -set for T' must contain one of $\{v_3, c\}$, we may assume it contains v_3 . Addition of $\{v_2, a\}$ to a $\gamma_c^{k-1}(T')$ -set now gives the contradiction $\gamma_c^k(T) \leq n - k - 2$.

Assume finally that v_3 has no leaf but besides v_2 and v_4 is adjacent to a_1, a_2, \dots, a_t , $t \geq 1$, where each a_i has valency two and is adjacent to the leaf b_i , $1 \leq i \leq t$.

We have $k - t \geq 1$ because $V(T) \setminus \{v_1, b_1, b_2, \dots, b_t, v_\alpha\}$ is a connected subgraph with $n - t - 2$ vertices which dominate T , so that $n - k - 1 = \gamma_c^k(T) \leq n - t - 2$ giving $k - t \geq 1$. Consider the tree $T' = T \setminus \{v_1, v_2, a_1, a_2, \dots, b_1, b_2, \dots, b_t, v_3\}$ of order $n - 2t - 3$.

If $n - 2t - 3 \geq 2(k - t) + 1$ we obtain by Proposition 3 that $\gamma_c^{k-t}(T') \leq (n - 2t - 3) - (k - t) - 1 \leq n - k - t - 4$, and by addition of the $t + 2$ vertices $\{v_2, v_3, a_1, a_2, \dots, a_t\}$, (which span a connected subgraph of T), to a $\gamma_c^{k-t}(T')$ -set we obtain $\gamma_c^k(T) \leq n - k - 2$, a contradiction. So we have $n - 2t - 3 \leq 2(k - t)$ and now $|V(T')| = n - 2t - 3 \leq 2(k - t)$ implies $\gamma(T') \leq \frac{|V(T')|}{2} \leq k - t$ which by remark 1 gives that $\gamma_c^{k-t}(T') = \gamma(T')$ and hence addition of the $t + 2$ vertices $\{v_2, v_3, a_1, a_2, \dots, a_t\}$ to a $\gamma_c^{k-t}(T')$ -set (having at most $k - t$ vertices) gives $\gamma_c^{k-t+1}(T) \leq k + 2$. We now have $n - k - 1 = \gamma_c^k(T) \leq \gamma_c^{k-t+1}(T) \leq k + 2$ giving $n \leq 2k + 3$, so the assumption $n \geq 2k + 3$ implies $n = 2k + 3$. By hypothesis $\gamma_c^k(T) = k + 2$ and we have $\gamma(T) \leq k + 1$ by Proposition 1.

Thus $\gamma(T) = k + 1$, (because otherwise $\gamma_c^k(T) = \gamma(T) < k + 2$), and any $\gamma(T)$ -set must consist of $k + 1$ isolated vertices. As $\gamma(T) = \lfloor \frac{n}{2} \rfloor$ the tree T by Proposition 4 is of type A or B. But T cannot be of type B, for assume T is of type B. Then T consists of a 3-path, uvw , with each of its ends joined to stems of corona trees, and since we have just seen that $v_3, v_{\alpha-2}$ are neither stems nor leaves, they must play the role of u, w , so $\alpha = 7$ and T consists of two subdivided stars centered respectively at $u = v_3$ and $w = v_5$ and a vertex $v = v_4$ joined to u and w . Among its γ -sets this tree T has one with two adjacent vertices, namely v_2 and v_3 , a contradiction, so T is of type A.

Using, in analogy to v_2, v_3 , that $d_T(v_{\alpha-1}) = 2$ and that $v_{\alpha-2}$ is not a stem, we get that $\alpha = 5$ and T is a subdivided star so that (iii) occurs.

Conversely, it is easy to see that if (i), (ii) or (iii) holds then $\gamma_c^k(T) = \gamma(T) = n - k + 1$. This proves Theorem 5. \square

References

- [1] X. Baogen, E.J. Cockayne, T.W. Haynes, S.T. Hedetniemi, Z. Shangchao: *Extremal graphs for inequalities involving domination parameters*, Discrete Math. 216 (2000), 1-10.
- [2] P. Duchet, H. Meyniel: *On Hadwiger's number and stability number*, Ann. Discr. Math. 13 (1982), 71-74.
- [3] J. F. Fink, M. S. Jacobson, L. F. Kinch, J. Roberts: *On graphs having domination number half their order*, Period. Math. Hungar. 16 (1985), 287-293.
- [4] S. Guha, S. Khuller: *Approximation algorithms for connected dominating sets*, Algorithmica 20 (1998), 374-387.
- [5] B.L. Hartnell, P.D. Vestergaard: *Partitions and domination in graphs*, J. of Combin. Math. and Combin. Comput. 46 (2003), 113-128.
- [6] B.L. Hartnell, P.D. Vestergaard: *Dominating sets with at most k components*, Ars Combin. 74 (2005), 223-229.
- [7] Teresa W. Haynes, Stephen T. Hedetniemi, Peter J. Slater: *Fundamentals of domination in graphs*. Marcel Dekker, New York 1998.
- [8] S.T. Hedetniemi, R.C. Laskar: *Connected domination in graphs*, B. Bollabas, ed.: *Graph Theory and Combinatorics*. Academic Press, London, 1984.
- [9] J. Niemen: *Two bounds for the domination number of a graph*, J. Inst. Math. Applics. 13 (1974), 183-187.
- [10] O. Ore: *Theory of Graphs*, Amer. Soc. Colloq. Publ. vol. 38. Amer. Math. Soc., Providence, RI 1962.
- [11] C. Payan and N. H. Xuong: *Domination-balanced graphs* J. of Graph Theory 6 (1982), 23-32.
- [12] B. Randerath, L. Volkmann, *Characterization of graphs with equal domination and covering number*, Discrete Math. 191 (1998), 159-169.
- [13] Zs. Tuza, P.D. Vestergaard: *Domination in partitioned graphs*. Discuss. Math. Graph Theory. 22 (2002), 199-210.
- [14] E. Sampathkumar: *Domination parameters of a graph* pp. 271-299 in Teresa W. Haynes, Stephen T. Hedetniemi, Peter J. Slater, eds.: *Domination in Graphs, Advanced Topics*. Marcel Dekker, New York 1998.
- [15] S. Seager: *Partition domination of graphs of minimum degree two*. Congr. Numer. 132 (1998), 85-91.

